

An interactive visualization approach to explore mass variations in the Earth system

Tobias Rawald, Mike Sips, Ingo Sasgen, and Doris Dransch

Abstract

We present an approach that supports the visual exploration of mass variations in the Earth system in space and time. In close collaboration with geo-scientists, we identified important requirements for our exploration tool. Using a combination of three-dimensional spheres, we enable them to compare the differences between simulated mass variations and the measured gravity field. Since these variations change over time, we developed a tool that allows to navigate the temporal scale. In addition, we supply a mechanism to dynamically activate/deactivate multiple components of the simulation model applied. These features enable scientists to better understand the impact of the individual processes on the measured Earth's gravity field. In this paper, we discuss in detail the geo-scientific requirements and how our tool addresses these requirements.

Categories and Subject Descriptors (according to ACM CCS): I.3.8 [Computer Graphics]: Applications —

1. Introduction

Satellite missions such as *GRACE* [Hel14, NAS14] provide high-precision measurements of the Earth's gravity field, which changes from one place to another depending upon the concentration of mass in a region. Mountains usually cause the gravitational force to be higher than it would be on a featureless planet. Conversely, the depression of the land-mass through the presence of glaciers million years ago usually causes the gravitational force to be lower. The Earth's gravity field also changes at finer time scales like effects associated with the hydrologic cycle. Analyzing the short-term changes in masses, geo-scientists gain a better understanding of the variability of the Earth's climate system, e.g., melting of ice caps.

Because of Newton's law of gravitation, a variety of masses usually influence the gravity signal at a particular region. The measured signal reflects the impact of several environmental processes, which we refer to as *subsystems*. This results in two challenges. The first challenge is to develop a better understanding of the measured gravity field. The interpretation of satellite measurements requires Earth system modelers to explore how different masses influence these measurements. This exploration involves the simultaneous comparison of the following four *dimensions*:

I The calculated mass variations based on a simulation model.

- II The inferred gravity field changes based on the simulated mass variations.
- III The satellite measurements of the Earth's gravity field changes.
- IV The inferred mass variations deduced from the satellite measurements.

The second challenge is to utilize the satellite measurements to improve simulation models capturing the short-term temporal dynamics of mass variations. This challenge requires Earth system modelers to explore the temporal variation of the gravity field and mass changes in different subsystems. Our tool addresses these challenges by providing four visual views; one for each dimension mentioned above (see Fig. 1). The corresponding visualization approach utilizes well-known techniques, such as color mapping, height mapping, and linked views.

These four views support the geo-scientists to relate mass variations to measured gravitational signals and therefore allow them to attribute mass variations to certain subsystems, e.g., attributing the high gravitational signal in Fig. 1 (III) to a high water level in the Amazon River (hydrological systems). The resulting tool supports the second challenge by providing a slider that allows them to explore mass variations over time. Additionally, Earth system modelers can utilize our tool to validate the geo-scientific algorithms applied.




Figure 1: The main window displays four spherical views; one for each dimension. The roman numerals are not part of the actual visualisation but rather serve as a visual aid to identify the dimension displayed. The temporal navigation is realized via a slider at the bottom. Additional information is displayed within a textual overlay.

2. Visualization and Interaction Concept

The data sets that reflect the mass variations and gravity field changes have specific spatial and temporal resolutions. The spatial resolution specifies the sampling size of the Earth's surface. We refer to a discrete sample of the Earth's surface as *grid*. In regard to our scenario, each grid has a resolution of a quarter of a degree concerning longitude as well as latitude. The temporal resolution specifies the number of time steps captured. Mass variations as well as gravity field changes are provided on a monthly basis. To address the properties of the data and the scientific challenges, in collaboration with geo-scientists we developed a visualization and interaction concept. We defined the following requirements our tool has to comply with:

1. Exploration of mass variation interdependencies,
2. Comparison of the different dimensions,
3. Exploration of mass variations and their impact to the gravity field in space and time, and
4. Exploration of mass variations in the individual subsystems.

To support these requirements, our interactive visualization tool has two major components. The first component is the *main window* that provides an overview over all four dimensions (see Fig. 1). For each dimension a three-

dimensional sphere visualization is included, depicting data referring to a certain month of a year. Our system supports the exploration in space and time by (a) allowing geo-scientists to rotate and zoom the spheres and (b) to choose a particular point in time via a slider. Note, the visualized data in each view has the same spatial resolution. To support the exploration of the impact of subsystems to the gravity field, our tool provides a *dialog window* for each of the four views (see Fig. 2). The dialog windows present visualization properties and allow the user to adjust the visualization, e.g., select new data by adding or removing subsystems. Each dialog window can be activated or deactivated individually.

In the upcoming sections, we discuss the visualization of the Earth system data, the requirements and the associated tasks in detail.

2.1. Exploration of mass variation interdependencies

Since a number of masses influence the measurement of the gravity field over a region, it is important that the visualization clearly presents these interdependencies. We choose to visualize the grid data using three-dimensional spheres. It allows to correlate mass variations on opposite sites of the globe easily. This requirement was explicitly given by the domain expert.




Figure 2: The dialog window allows the user to activate or deactivate subsystems in the visualization. It allows to switch the display mode and presents the color mapping.

Visualizing the grid data in three-dimensional space introduces visibility problems; namely, not all parts of a sphere are visible to the user simultaneously. Therefore, our tool provides means of navigation to focus on a region of interest. The user can manipulate the position of the camera, which creates the impression of rotating the sphere along the X and Y axis or to zoom-in and zoom-out. Since an important task is to support geo-scientists to compare variations across multiple dimensions, our system synchronizes the user interaction across the four spheres. The Earth system modeler can interact with the visualization by pressing the arrow keys for rotating the spheres as well as the "+" and the "-" key to zoom-in and zoom-out.

2.2. Comparison of different dimensions

Regarding the comparisons of dimensions, we distinguish the following tasks:

- (I vs. II) Compare the simulated mass variations with the inferred Earth's gravity field changes.
- (III vs. IV) Compare the measured gravity field changes with the inferred mass variations.
- (I vs. IV) Compare the simulated with the inferred mass variations.
- (II vs. III) Compare the inferred with the measured gravity field changes.

We support the comparison of dimensions by arranging the spheres in a special layout; each horizontal and vertical adjacent pair of spheres refers to one of the four tasks (see Fig. 1). Additionally, each view presents further information within a textual overlay, e.g., the type of the dimension displayed.

2.3. Exploration of mass variations in space and time

To identify interesting patterns in space and time, the geo-scientist has to:

- Navigate through time to identify potentially interesting mass variations.
- Identify and focus on a region with unusual mass variations.
- Distinguish between local peaks and large-area variations.
- Connect the interesting region to its geographical location.

To support the first task, our tool synchronizes the four spheres concerning the temporal navigation. All four views present data referring to the same point in time. The user can interactively navigate through time via a slider on the bottom of the main window. Each tick of the slider represents a specific month of a year.

To support geo-scientists to identify interesting mass variations, our tool utilizes a combination of *color* and *size* to visually encode mass variations and changes in the gravity field [Ber11]. In the following, we refer to size as the distance between the center of the sphere and a point on its surface.

The challenge in utilizing *color* as retinal variable is that mass variations and gravity field changes differ in their domain. We address this challenge in our tool by applying different value range intervals for mass variations (I and IV) and gravity field changes (II and III). Note, the same color palette is applied in all four views. The mapping to color is displayed in the corresponding dialog window for each dimension (see Fig. 2). We incorporate the domain knowledge of Earth system modelers to create meaningful visual representations of the data. An important convention is to use bluish colors to encode negative variations and reddish colors to encode positive changes. To address this convention, we use the color scheme *11-class RdBu* provided by *Color Brewer 2.0* [BHT14]. Note, the center of the scheme is a color close to white, which in our case operates as neutral value, meaning that no variation is present.

To further support the distinction between local peaks and large-area variations, we utilize the second retinal variable *size*. We enhance the visual encoding by varying the spheres radius for each location based on the associated mass or gravity field variation. Again, we define a default radius that represents the neutral value. A positive variation leads to a radius greater and a negative variation to a radius smaller

than the default radius. As a result, the spheres appear distorted. Our color encoding in combination with the distortion supports Earth system modelers to identify interesting mass variations. The mapping to size simplifies the distinction between local peaks and large-area variations. To support effective exploration, the user can switch between two display modes. The first employs only color to encode the variations (*Plane*), whereas the second combines the encoding using color and size (*Elevation*).

The fourth important task is to relate unusual mass variations to geographic regions. To support this task, our sphere visualization presents visual landmarks. This comprises a regular grid of prominent longitudes and latitudes as well as the coastlines of continents on the sphere. To distinguish the visual landmarks from the mass variation data, we assign colors that differ from the colors used to visualize the grid data. Additionally, we also adapt the visual landmarks in the display mode *Elevation* according to the radius computed for every grid point.

2.4. Exploration of the individual subsystems

This requirement is a central aspect of understanding the impact of a single subsystem to the measured gravity signal. Geo-scientists determine its influence by comparing the intensity of a subsystem to the measured gravity field or the inferred mass variation. Our tool enables to attribute the high gravitational signal in Fig. 1(III) to the hydrological subsystem. The hydrological subsystem shows a high peak at the Amazon River which resembles the measured gravity signal. The activation of other subsystems, e.g., in Fig. 1(II) the subsystem *glacial-isostatic adjustment* (GIA) is also activated, did not change the simulated gravity signal.

Hence, our tool has to support the following two tasks:

- Activate/deactivate a subsystem signal belonging to the simulated mass change variation dimension.
- Activate/deactivate a subsystem signal belonging to the inferred gravity field change dimension.

The user can activate/deactivate subsystems in the dimensions (I) and (II). We provide controls to support this analytical step within the dialog window (see Fig. 2). Each subsystem signal can either be *activated* or *deactivated*. Therefore, we supply a binary check-box for each subsystem available in the data. After subsystems are activated or deactivated, our tool updates the corresponding sphere visualization automatically.

3. Related Work

Prior to our own efforts, we evaluated the *World Wind SDK* that is provided by NASA [Nat11]. We assessed its performance in displaying our high-resolution grid data, each sphere comprises over 10^6 data points, as not sufficient.

The visualization and interactive exploration require the

integration of measured and simulated data into a common data model. The simulation data is generated by independent or coupled physical models of the subsystems of the Earth; here, it includes continental hydrology, cryosphere, ocean, and GIA. To support exploratory analysis, our visualization system needs to have efficient access to spatial (grid points) and temporal data (time steps). We developed an efficient data model to enable fast access to the spatial and temporal data [SRSU12]. In contrast to [SRSU12], we here describe the comprehensive task analysis and design rationale for our visual exploration tool.

The comparison of simulated and measured data is an important challenge in many geo-scientific fields. It has stimulated research projects in the visualization community, e.g., [USKD12], to exploit the benefits of interactive visualization for this comparison. Geo-scientists use visualization and exploration tools to extract information and meaning from complex geo-scientific data (see for example [Eur13]). Commercial systems such as Avizio [FEI14] and scientific systems like NCAR Graphics [Nat14] are available to support the exploration of complex geo-scientific data sets.

4. Conclusion

In cooperation with Earth system modelers, we developed a tool that allows to visually explore spatial and temporal mass variations and their impact to the Earth's gravity field. Our tool supports the central task of comparing the measured gravity field with simulated mass variations. To develop a visualization and interaction concept, we regularly met with geo-scientists to identify the spot at which interactive visualization can provide a valuable contribution to the scientific problem. On this basis, we developed requirements for an exploration tool. We iteratively refined these requirements to identify the underlying analytical tasks. Finally, we derived appropriate visual metaphors and interaction mechanisms to support interactive visual analysis and exploration.

The key features of our tool are three-dimensional visualizations of mass variations and the gravity field changes. Our tool provides means of navigation in space and time to identify and focus on interesting variations. Geo-scientists use our prototypical exploration tool in multiple scientific scenarios, e.g., to analyze incoming data from current satellite missions. A next step is to integrate our exploration tool in a central system environment for the management of Earth system data. For this purpose, we need to connect the tool to additional databases. Moreover, we plan to extend the visualization methods to support temporal analysis through slicing the spheres; each slice representing a different point in time. Concatenating the slices, temporal navigation is implemented by rotating spheres. This approach would make the slider obsolete.

References

- [Ber11] BERTIN J.: *Semiology of Graphics: Diagrams, Networks, Maps*. Esri Press, Redlands, 2011. 3
- [BHT14] BREWER C., HARROWER M., THE PENNSYLVANIA STATE UNIVERSITY: ColorBrewer: Color Advice for Maps, March 2014. URL: <http://www.colorbrewer2.org/>. 3
- [Eur13] EUROPEAN GEOSCIENCES UNION GENERAL ASSEMBLY 2013: Techniques and tools for effective visualization in the geosciences, March 2013. URL: <http://meetingorganizer.copernicus.org/EGU2013/orals/12093>. 4
- [FEI14] FEI VISUALIZATION SCIENCES GROUP: Avizio Visualization Software, March 2014. URL: <http://www.vsg3d.com/avizo/overview>. 4
- [Hel14] HELMHOLTZ CENTRE POTSDAM - GFZ GERMAN RESEARCH CENTRE FOR GEOSCIENCES: Gravity Recovery and Climate Experiment (GRACE) Mission, March 2014. URL: <http://www.gfz-potsdam.de/en/research/organizational-units/departments-of-the-gfz/department-1/global-geomonitoring-and-gravity-field/topics/development-operation-and-analysis-of-gravity-field-satellite-missions/grace/>. 1
- [NAS14] NASA EARTH OBSERVATORY: Gravity Recovery and Climate Experiment (GRACE), March 2014. URL: <http://earthobservatory.nasa.gov/Features/GRACE/>. 1
- [Nat11] NATIONAL AERONAUTICS AND SPACE ADMINISTRATION: World Wind Java SDK, July 2011. URL: <http://worldwind.arc.nasa.gov/java/>. 4
- [Nat14] NATIONAL CENTER FOR ATMOSPHERIC RESEARCH: NCAR Graphics, March 2014. URL: <http://ngwww.ucar.edu/>. 4
- [SRSU12] SIPS M., RAWALD T., SASGEN I., UNGER A.: Exploration of spatial and temporal signatures in multiple components of a geodetic earth system model. In *Managing Resources of a Limited Planet: Pathways and Visions under Uncertainty*, Sepelt R., Voinov A., Lange S., Bankamp D., (Eds.). 2012. 4
- [USKD12] UNGER A., SCHULTE S., KLEMANN V., DRANSCH D.: A visual analysis concept for the validation of geoscientific simulation models. *IEEE Transactions on Visualization and Computer Graphics* 18, 12 (2012), 2216 – 2225. 4